



3 1799 00013 5087

ACQUISITION

MAY 24 1990

COLORADO STATE LIBRARY
State Publications Library

Colorado
State
University
Cooperative
Extension

no. 4.908

service in ACTION

Evaluation of drinking water quality for livestock

P. N. Soltanpour and W. L. Raley¹

Quick Facts

Livestock drinking water should be tested for salinity and toxic elements if water quality is not known.

A "representative" water sample should be taken for any testing.

The National Academy of Sciences has proposed general guidelines for use of saline waters and for upper limits of toxic ions in water.


There are no easy answers or quick fixes for toxic water problems.

Excessive salinity in drinking water that is used for livestock can upset water balance of livestock and sometimes cause animal deaths. The National Academy of Sciences has proposed guidelines for use of saline waters for livestock and poultry (Table 2).

High levels of specific ions in water can cause animal health problems and death. The National Academy of Sciences has proposed upper limits for toxic substances in water (Table 1).

Unsafe levels of salts and ions depend on the amount of water consumed each day and the weight of the animal. Livestock drinking water should be tested for salinity and toxic elements if water quality is not known. The general guidelines presented in tables 1 and 2 include an appropriate margin of safety. For a more specific interpretation of drinking water quality for livestock, contact your veterinarian.

To obtain a "representative" water sample for testing, collect approximately 1-pint samples at random intervals using clean glass or plastic containers. Pour small samples into large con-



tainer and thoroughly mix. Take final sample from mixed large container. Avoid taking sample from inactive wells or directly after drilling. Thoroughly pump or bail wells before sampling.

If a water test report shows that water contains toxic substances that exceed the upper limit guidelines listed in tables 1 and 2, treatment may be necessary before livestock drink the water. However, there usually are no simple answers or quick fixes for toxic water problems. In many cases, water treatment may be impractical or too costly. Some circumstances may warrant the use of *ion-exchange filters*, *distillation* or *dilution* to correct the problem. Consultation with a sanitary engineer or water treatment specialist on a case-by-case basis is the best recommendation.

¹P. N. Soltanpour, Colorado State University Cooperative Extension agronomy specialist and professor, agronomy; and W. L. Raley, Cooperative Extension Western District director (8/89)

Table 1: Recommendations for levels of toxic substances in drinking water for livestock.

Constituent	Upper limit
Aluminum (Al)	5 mg/l
Arsenic (As)	0.2 mg/l
Beryllium (Be)	no data
Boron (B)	5.0 mg/l
Cadmium (Cd)	.05 mg/l
Chromium (Cr)	1.0 mg/l
Cobalt (Co)	1.0 mg/l
Copper (Cu)	0.5 mg/l
Fluoride (F)	2.0 mg/l
Iron (Fe)	no data
Lead (Pb)	0.1 mg/l ¹
Manganese (Mn)	no data
Mercury (Hg)	.01 mg/l
Molybdenum (Mo)	no data
Nitrate + nitrite (NO ₃ - N, NO ₂ -N)	100 mg/l
Nitrite (NO ₂ -N)	10 mg/l
Selenium (Se)	0.05 mg/l
Vandium (V)	0.10 mg/l
Zinc (Zn)	24 mg/l
Total dissolved solids (TDS)	10,000 mg/l ²

¹Lead is accumulative and problems may begin at threshold value = 0.05 mg/l.

²See Table 2.

Sources: Environmental Studies Board, Nat. Acad. of Sci., Nat. Acad. of Eng., **Water Quality Criteria**, 1972.

Ayers, R.S. and D.W. Westcot. **Water Quality for Agriculture**. Food and Agriculture Organization of the United Nations, Rome, 1976.

Table 2: Guide to the use of saline waters for livestock and poultry.

Total soluble salts Content of waters	
Less than 1000 mg/l (EC < 1.5 mmhos/cm)	Relatively low level of salinity. Excellent for all classes of livestock and poultry.
1000 - 3000 mg/l (EC = 1.5 - 5 mmhos/cm)	Very satisfactory for all classes of livestock and poultry. May cause temporary and mild diarrhea in livestock not accustomed to them; may cause watery droppings in poultry.
3000 - 5000 mg/l (EC = 5 - 8 mmhos/cm)	Satisfactory for livestock, but may cause temporary diarrhea or be refused at first by animals not accustomed to them. Poor waters for poultry, often causing watery feces, increased mortality and decreased growth, especially in turkeys.
5000 - 7000 mg/l (EC = 8 - 11 mmhos/cm)	Can be used with reasonable safety for dairy and beef cattle, sheep, swine and horses. Avoid use for pregnant or lactating animals. Not acceptable for poultry.
7000 - 10,000 mg/l (EC = 11 - 16 mmhos/cm)	Unfit for poultry and probably for swine. Considerable risk in using for pregnant or lactating cows, horses or sheep, or for the young of these species. In general, use should be avoided although older ruminants, horses, poultry and swine may subsist on them under certain conditions.
Over 10,000 mg/l (EC > 16 mmhos/cm)	Risks with these highly saline waters are so great that they cannot be recommended for use under any condition.

Sources: Environmental Studies Board, Nat. Acad. of Sci., Nat. Acad. of Eng., **Water Quality Criteria**, 1972.

Ayers, R.S. and D.W. Westcot. **Water Quality for Agriculture**. Food and Agriculture Organization of the United Nations, Rome, 1976.